

## WELL CHILD EXAM - EARLY CHILDHOOD 2 YEARS

7311	DEPARTMENT OF SOCIAL 6-HEALTH SERVICES	(Meets EPSDT Guidelines)				DATE	
		EARLY C	HILDHO	OD: 2 YEARS			
	CHILD'S NAME		BROU	GHT IN BY	DATE	OF BIRTH	
PARENT TO COMPLETE ABOUT THE CHILD	ALLERGIES	RGIES CURRENT MEDICATIONS					
	ILLNESSES/ACC	DENTS/PROBLEMS/CONCERNS SINC	E LAST V	/ISIT	TODAY I HAVE A QUEST	ION ABOUT:	
	□ □ My	child eats a variety of foods. child settles well at bedtime and sle	eeps wel		y child can stack blocks. y child uses 2-3 word ser y child is showing interes		
VEIGHT KG./OZ. PERCENTILE HEIGHT CM/IN. PERCENTILE HEAD CIR. PERCENTILE				Diet			
Review of systems Review of family history  Screening:  N A Hearing Vision Development: Circle area of concern				Elimination  Sleep  Review Immunization Record Lead Exposure  Fluoride Supplements Fluoride Varnish  Hct/Hgb TB Dental Referral  Health Education: (Check all discussed/handouts given)  Nutrition Toilet Training Safety  Development Car Seat or Booster Seat TV Habits  Passive Smoking Discipline/Limits Teeth Care			
Adaptive/Cognitive Language/Communication  Gross Motor Social/Emotional Fine Motor  Behavior							
Mental	Mental Health		☐ Child Care ☐ Other:				
Skin Head Eyes C Ears Nose Oropha Neck Nodes	al appearance Cover/Uncover arynx/Teeth	N A Chest Lungs Cardiovascular/Pulses Abdomen Genitalia Spine Extremities Neurologic Gait ngs and comments.	N A	Assessment/Plan			
				IMMUNIZATIONS GIV	/EN		
				REFERRALS			
NEXT VISIT: 3 YEARS OF AGE				HEALTH PROVIDER NAME			
4EALTH PROVIDER SIGNATURE				HEALTH PROVIDER ADDRESS			

DSHS 13-684A(X) (REV. 08/2004) - TRANSLATED

 $\square$  SEE DICTATION

DISTRIBUTION: White - Physician Yellow - Parent/Guardian

## Your Toddler's Health at 2 Years

## **Milestones**

# Ways your toddler is developing between 2 and 3 years of age.

Jumps in place.

Pedals a tricycle.

Speaks in three or four word sentences.

Feeds and dresses himself.

Can draw a cross and a circle with a crayon.

Plays "make believe" games with dolls and stuffed animals.

You help your child learn new skills by playing with her.

## For Help or More Information

## Safe Gun Storage Information: 1-800-LOK-IT-UP (565-4887)

**Child Care:** Washington State Child Care Resources & referral Network,

1-800-446-1114

**Poisoning Prevention:** The Washington State Poison Center, 1-800-732-6985 or 1-800-572-0638 (TTY Relay)

For help if you feel very frustrated with your child: Family Help line, 1-800-932-HOPE (4673) Local crisis hotlines.

Parenting Skills or Support: Family Help line, 1-800-932-HOPE (4673) Family Resources Northwest, 1-888-746-9568 Local Community College Classes

## **Health Tips**

Are your child's immunizations up to date? If so, he will not need any more until just before he enters kindergarten.

Offer your child a variety of healthy foods every day. Limit junk foods. Eat meals together as a family as often as possible. Turn off the TV while eating together.

Brush your child's teeth at least once a day with a pea-sized amount of fluoride toothpaste. Make sure she gets a dental checkup once a year.

## **Parenting Tips**

Talk with your child about what he is doing. Read picture books together and talk about the pictures.

Help your child enjoy active games like catch, tag, and hide-and-seek. Give her simple toys to play with, like blocks, crayons and paper, stuffed animals.

You may want your child to be toilet trained soon, but he may not be ready until about age 3. He will show you he is ready by being dry after sleep and telling you when he wants to use the toilet.

Limit television to 1 hour or less for children. Watch with them and talk about the programs.

## **Safety Tips**

- Keep cleaning supplies and medicines locked up out of reach.
- Always hold your child's hand while walking near traffic, including in parking lots. Check behind your car before backing up, in case a child is behind it.
- If you have guns at home, always store them safely. Keep them unloaded and locked up.
- Put a life jacket on your child whenever she is near the water or in a boat. Always watch her around water.
- Keep matches and lighters out of reach.

## Guidance to Physicians and Nurse Practitioners for Early Childhood (2 years)

The following highlight EPSDT screens where practitioners often have questions. They are not comprehensive guidelines.

#### Fluoride Screen

Fluoride supplements should be considered for all children drinking fluoride deficient (<0.6 ppm F) water. Before supplements are prescribed, it is essential to know the fluoride concentration of the patient's drinking water. Once the fluoride level of the water supply has been evaluated, either through contacting the public health officials or water analysis, as well as evaluating other sources of fluoride, the daily dosage schedule can be recommended. Pediatric Dentistry: Reference Manual 1999--00.(21)5.

#### **Lead Screen**

Screen children for these risk factors:

- Live in or frequently visits day care center, preschool, baby sitter's home or other structure built before 1950 that is dilapidated or being renovated.
- Come in contact with other children with known lead toxicity (i.e., blood lead 15 ug/dl).
- Live near a lead processing plant or with parents or household members who work in a lead-related occupation (e.g., battery recycling plant).

## **Pneumococcal Screen**

Screen children for these risk factors:

 Age 2 and over with chronic illnesses specifically associated with pneumococcal disease or its complications; anatomic or functional aspenia; sickle cell disease; nephritic syndrome or chronic renal failure; cerebrospinal fluid leaks; or conditions associated with immunosuppression.

## **Hepatitis B Vaccine**

• For children and adolescents not vaccinated against hepatitis B in infancy, begin the hepatitis B vaccine series during any childhood visit. Give the second dose at least one month after the first dose and give the third dose at least four months after the first dose and at least two months after the second dose.

## **Developmental Milestones**

Always ask about and follow-up on parent concerns about development or behavior. You may use the following screening list, or use the Ages and Stages Questionnaire, the Denver II, the ELMS2 (a language screen), or the MacArthur Communication Development Inventory.

Yes	No	
		Kicks ball forward.
		Removes article of clothing (not hat).
		Combines 2 words.
		Uses own name to refer to self.
		Vocabulary of more than 50 words.
		Strangers understand half child's speech.
		Understands a two-step verbal command ("Pick up the toy; put it away") without gestures.
		Points to 6 named body parts (nose, eyes, ears, mouth, hands, feet, tummy, hair).
		Stacks 4 blocks.

Persistent rocking, hand flapping, head banging, or toe walking.

• No spontaneous two word (not echolalic) phrases.

· Any loss of any language or social skills.

Instructions for developmental milestones: At least 90% of children should achieve the underlined milestones by this age. If you have checked "no" on any two items, even one of the underlined items, or any of the boxed items (abnormal behavior at this age), refer the child for a formal developmental assessment.

Notes: Immunization schedules are from the Advisory Committee on Immunization Practice of the U.S. Centers for Disease Control and Prevention. Parents and providers may call Healthy Mothers, Healthy Babies (1-800-322-2588) with questions or concerns on childhood development.